

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 51/16		A1	(11) International Publication Number: WO 99/52849
			(43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/US99/07466 (22) International Filing Date: 5 April 1999 (05.04.99) (30) Priority Data: 60/081,202 9 April 1998 (09.04.98) US 9810188.4 13 May 1998 (13.05.98) GB (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LI, Jing [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). TSCHAEN, David, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). SONG, Zhiguo [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ZHAO, Mangzu [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: OXIDATION PROCESS USING TEMPO			
(57) Abstract			
The present invention relates to the oxidation of a primary alcohol of Formula (II) to the carboxylic acid of Formula (I): $R^1CH_2OH \rightarrow R^1CO_2H.$			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION
OXIDATION PROCESS USING TEMPO

BACKGROUND OF THE INVENTION


5 Oxidation is one of the most fundamental transformations
in organic synthesis and there are numerous methods reported in the
literature. (Hudlicky, M. "Oxidations In Organic Chemistry", ACS
Monograph No. 186 American Chemical Society Washington D.C.
(1990).) However, relatively few methods exist for the oxidation of
10 primary alcohols to the corresponding carboxylic acids. The most
commonly used ones are $\text{CrO}_3/\text{H}_2\text{SO}_4$ (Bowden; Heilbron; Jones; Weedon
J. Chem. Soc., 1946, 39; Bowers; H.; Jones; L. *J. Chem. Soc.*, 1953, 2548;
Millar, J. G.; Oehlschlager, A. C.; Wong, J. W. *J. Org. Chem.* 1983, 48,
4404.), $\text{RuCl}_3/\text{H}_5\text{IO}_6$ (Carlsen, P. H. J.; Katsuki, T.; Martin V. S.;
15 Sharpless, K. B. *J. Org. Chem.* 1981, 46, 3936.) and TEMPO/ NaClO
(Nooy, A. E. J. de; Besemer, A. C.; Bekkum, H. v. *Synthesis*, 1996, 1153.;
Anelli, P. L.; Biffi, C.; Montanari, F.; Quici, S. *J. Org. Chem.* 1987, 52,
2559.; Miyazawa, T.; Endo, T.; Shiihashi, S.; Okawara, M. *J. Org.*
Chem. 1985, 50, 1332). A two-step process involving Swern oxidation
20 (Mancuso, A. J.; Huang, S-L., Swern, D. *J. Org. Chem.* 1978, 43, 2480.;
Mancuso, A. J.; Brownfan, D. S.; Swern, D. *J. Org. Chem.* 1979, 44,
4148.; Ireland, R.; Norbeck, D. *J. Org. Chem.* 1985, 50, 2198.) followed by
oxidation of the resulting aldehyde with NaClO_2 (Lindgren, B. O.;
Nilsson, T. *Acta Chem. Scand.* 1973, 27, 888.; Dalcanele, E.; Montanari,
25 F. *J. Org. Chem.* 1986, 51, 567) is another option. All of these procedures
have some limitations and disadvantages and new methods for the
oxidation of primary alcohols to the carboxylic acids are still desired.
(Schroder, M.; Griffith, W. P. *J. Chem. Soc. Chem. Comm.* 1979, 58.;
and Paquette, L. A.; Dressel, J.; Pansegrau, P. D. *Tetrahedron Lett.*
30 1987, 28, 4965.)

 The present invention relates to an oxidation using sodium
chlorite in the presence of a catalytic amount of TEMPO and sodium
hypochlorite which converts a primary alcohol to a carboxylic acid. This
oxidation method avoids the disposal issues associated with running a
35 Jones oxidation ($\text{CrO}_3/\text{H}_2\text{SO}_4$) reaction, as well as reducing the

epimerization of any α -chiral centers and is a one step procedure. For substrates prone to chlorination with the TEMPO-NaClO protocol, the present invention reduces this problem.

5 SUMMARY OF THE INVENTION

The present invention discloses a process for preparing a compound of Formula I:



wherein:

10 R^1 is:

- a) H,
- b) C₁-C₈ alkyl,
- c) C₂-C₈ alkynyl,
- d) C₃-C₇ cycloalkyl,
- 15 e) aryl,
- f) heteroaryl, or
- g) heterocyclyl;

C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl, are unsubstituted or substituted with one, two or three
20 substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₃-C₈ cycloalkyl, aryl, heteroaryl, heterocyclyl, and CO(CH₂)_nCH₃,

aryl is defined as phenyl or naphthyl, which is unsubstituted or substituted with one, two or three substituents selected from
25 the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, or when aryl is substituted on adjacent carbons they can form a 5- or 6-membered fused ring
30 having one, two or three heteroatoms selected from O, N, and S, this ring is unsubstituted or substituted on carbon or nitrogen with one, two or three substituents selected from

the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, and CO(CH₂)_nCH₃;

heteroaryl is defined as a 5- or 6-membered aromatic ring
5 containing 1, 2 or 3 heteroatoms selected from O, N and S ,
which is unsubstituted or substituted with one, two or three
substituents selected from the group consisting of: OH,
CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈
alkynyl, C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and additionally
10 the 5- or 6-membered aromatic ring can be benzofused and
unsubstituted or substituted with one, two or three
substituents as described above;

heterocyclyl is defined as a 5- or 6-membered, non-aromatic ring
containing 1, 2 or 3 heteroatoms selected from O, N and S ,
15 which may contain one or two double bonds and which is
unsubstituted or substituted with one, two or three
substituents selected from the group consisting of: OH,
CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈
alkynyl, or C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and
20 additionally the 5- or 6-membered ring can be benzofused
and unsubstituted or substituted with one, two or three
substituents as described above;

n is: 0 to 5;


25

t is: 0, 1 or 2;

R⁴ is: H, or C₁-C₈ alkyl; or

30 comprising the following steps:

1) adding to a compound of Formula II in a solvent,




a solution of phosphate buffer to maintain a pH of about 4.0 —
to about 8.0;

- 5 2) maintaining the phosphate-buffered biphasic mixture of the
 compound of Formula II at about 0°C to about 50°C;
- 3) adding a catalytic amount of TEMPO to the mixture; and
- 10 4) charging the TEMPO/phosphate-buffered biphasic mixture
 with a solution of sodium chlorite and a catalytic amount of
 sodium hypochlorite to oxidize to the compound of Formula I.

DETAILED DESCRIPTION OF THE INVENTION

- 15 The present invention discloses a process for preparing a
 compound of Formula I:



wherein:

R¹ is:

- 20 a) H,
 b) C₁-C₈ alkyl,
 c) C₂-C₈ alkynyl,
 d) C₃-C₇ cycloalkyl,
 e) aryl,
25 f) heteroaryl, or
 g) heterocyclyl;

C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl, are
unsubstituted or substituted with one, two or three
substituents selected from the group consisting of: OH,

CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₃-C₈ cycloalkyl, aryl, heteroaryl, heterocyclyl, and CO(CH₂)_nCH₃,

5 aryl is defined as phenyl or naphthyl, which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, or when aryl is substituted on adjacent
10 carbons they can form a 5- or 6-membered fused ring having one, two or three heteroatoms selected from O, N, and S, this ring is unsubstituted or substituted on carbon or nitrogen with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, and CO(CH₂)_nCH₃;

15 heteroaryl is defined as a 5- or 6-membered aromatic ring containing 1, 2 or 3 heteroatoms selected from O, N and S, which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and additionally
20 the 5- or 6-membered aromatic ring can be benzofused and unsubstituted or substituted with one, two or three substituents as described above;

25 heterocyclyl is defined as a 5- or 6-membered, non-aromatic ring containing 1, 2 or 3 heteroatoms selected from O, N and S, which may contain one or two double bonds and which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and additionally
30 the 5- or 6-membered ring can be benzofused and

unsubstituted or substituted with one, two or three substituents as described above;


n is: 0 to 5;

t is: 0, 1 or 2;

R⁴ is: H, or C₁-C₈ alkyl; or

comprising the following steps:

1) adding to a compound of Formula II in a solvent,



a solution of phosphate buffer to maintain a pH of about 4.0 to about 8.0;

2) maintaining the phosphate-buffered biphasic mixture of the compound of Formula II at about 0°C to about 50°C;

3) adding a catalytic amount of TEMPO to the mixture; and

4) charging the TEMPO/phosphate-buffered biphasic mixture with a solution of sodium chlorite and a catalytic amount of sodium hypochlorite to oxidize to the compound of Formula I.

The process as recited above, wherein the solvent is selected from the group consisting of: acetonitrile, tetrahydrofuran, acetone, tertiary C₄-C₈-alcohol, diethyl ether, DME (dimethyl ether), diglyme, triglyme, MTBE (methyl t-butyl ether), toluene, benzene, hexane, pentane, dioxane, dichloromethane, chloroform, carbon tetrachloride, or a mixture of said solvents.

The process as recited above, wherein the phosphate buffer comprises an aqueous mixture of NaOH, KOH, NaH₂PO₄, KH₂PO₄,

Na_2HPO_4 , and K_2HPO_4 , sufficient to maintain a pH of about 4.0 to about 8.0, and preferably a pH of about 6.5 to about 7.0.

The process as recited above, wherein TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy, free radical) is used in about 1.0 to about 10.0 mole percent, preferably about 5.0 to about 7.0 mole percent.

The process as recited above, wherein sodium chlorite is used in about 1.0 to about 3.0 equivalents, and preferably about 2.0 equivalents relative to the compound of Formula II.

The process as recited above, wherein sodium hypochlorite is used in about 1.0 to about 7.0 mole percent, preferably about 2.0 to about 5.0 mole percent.

The process as recited above, wherein the reaction temperature is about 0°C to about 50°C , and preferably about 35°C to about 40°C .

The process as recited above, wherein the reaction time is up to about 24 hours, and preferably between about 2 and about 4 hours.

It is further understood that the substituents recited above would include the definitions recited below.

The alkyl substituents recited above denote straight and branched chain hydrocarbons of the length specified such as methyl, ethyl, isopropyl, isobutyl, tert-butyl, neopentyl, isopentyl, etc.

Cycloalkyl denotes rings composed of 3 to 8 methylene groups, each of which may be substituted or unsubstituted with other hydrocarbon substituents, and include for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and 4-methylcyclohexyl.

The alkoxy substituent represents an alkyl group as described above attached through an oxygen bridge.


The aryl substituent represents phenyl and 1-naphthyl or 2-naphthyl, including aryls substituted with a 5- or 6-membered fused ring, such as an unsubstituted and substituted methylenedioxy, oxazolyl, imidazolyl, or thiazolyl ring.

The heteroaryl substituent represents a carbazolyl, furanyl, thienyl, pyrrolyl, isothiazolyl, imidazolyl, isoxazolyl, thiazolyl, oxazolyl, pyrazolyl, pyrazinyl, pyridyl, pyrimidyl, purinyl.

The heterocyclyl substituent represents, oxazolidinyl, thiazolidinyl, thiazolidinyl, oxadiazolyl, or thiadiazolyl.

Each of the above substituents (alkyl, alkynyl, alkoxy, cycloalkyl, aryl, heteroaryl, and heterocyclyl) can be either
5 unsubstituted or substituted as defined within the description.

Recently, in an attempt to oxidize primary alcohols, such as 1m:




1m

to the corresponding carboxylic acid, we found that $\text{RuCl}_3/\text{H}_5\text{IO}_6$ protocol offered low yield of the desired products. See Carlsen, P. H. et al. *J. Org. Chem.* **1981**, *46*, 3936. It was probably due to the destruction of electron rich aromatic ring. TEMPO catalyzed oxidation with bleach also gave low yield due to significant chlorination of the aromatic rings. See A. E. J. de Nooy, et al. *Synthesis*, **1996**, 1153.; P. L. Anelli, et al. S. *J. Org. Chem.* **1987**, *52*, 2559. and T. Miyazawa, et al. *J. Org. Chem.* **1985**, *50*, 1332. The synthesis of 1m is described in Merck Case No. 20127PV, entitled "Oxidation Process Using TEMPO" which is being filed simultaneously with this application.

In order to eliminate the chlorination problem, a few other oxidants (H_2O_2 , AcO_2H , $t\text{-BuO}_2\text{H}$ etc.) were examined, but no satisfactory results were obtained. Finally, when sodium chlorite (NaClO_2) was used as the oxidant, the product were obtained in 70-90% yield. The reaction appeared to be very slow (1-2%/hour) but generally went to completion overnight (~20 hours). More careful monitoring of the reaction indicated

that it was self accelerating process e.g. the conversion was less than 5% after one hour but reached ~90% in only 6 hours. Apparently, some more active species was generated as the reaction progress. Sodium hypochlorite (NaClO, bleach) was believed to be the most likely
5 candidate. Indeed, when 10mol% of bleach was added to the reaction mixture, the reaction was accelerate dramatically. It reached >50% conversion in one hour and finished in approximately three hours.

SCHEME 1




10

The reaction was then optimized regard to further reduce the chlorination and enhance the safety for scale up. The reaction was faster at lower pH, but it was accompanied by increased chlorination. It was slower at lower temperature as expected, but surprisingly, the
15 chlorination level appeared to be slightly elevated. Increasing the amount of TEMPO and bleach increased the reaction rate, but the TEMPO/NaClO ratio should be >2 to reduce the chances of chlorination. The bleach was added slowly and simultaneously with NaClO₂ to the batch at 35 °C to prevent build up of the oxidant and the risk of a run
20 away reaction. It should be noted that mixing of bleach and NaClO₂ prior to the addition is not advised since some toxic and potentially explosive chlorine dioxide (ClO₂) may be generated.

Next, a number of primary alcohols were oxidized to the carboxylic acids and the results are summarized in Table 1. In general,
25 the reaction were very smooth and the yield were excellent (85-100%).

Chiral alcohols **1g**, **1j**, and **1k** were oxidized to the corresponding carboxylic acid without any racemization of the labile chiral centers.

Mostly notably, for substrates prone to chlorination (**1c-1h**), our new procedure gave much better yields. The most dramatic demonstration of the superiority of our new procedure was revealed in entry 5. When **1e** was treated with NaClO and catalytic TEMPO, the desired product was obtained in less than 5% yield. One of the major side product was isolated and identified to be the chlorinated compound **4**,



based on NMR studies. On the other hand, our TEMPO/NaClO₂ protocol offered essentially quantitative yield of **2e**.

Table 1: TEMPO Catalyzed Oxidation of Primary Alcohols to Carboxylic Acids

5

























Substrate	Product	Yield (NaClO ₂)	Yield (NaClO)
 1a	 2a	98%	-
 1b	 2b	100%	-
 1c	 2c	99%	65%
 1d	 2d	100%	86%
 1e	 2e	96%	<5%
 1f	 2f	96%	80%

Table 1: (Cont.) TEMPO Catalyzed Oxidation of Primary Alcohols to Carboxylic Acids

5

Substrate	Product	Yield (NaClO ₂)	Yield (NaClO)
 1g	 2g	92%	60%
 1h	 2h	90%	20%
 1i	 2i	95%	-
 1j	 2j	95%	-
 1k	 2k	85%	-
 1l	 2l	100%	-

In conclusion, an efficient and environmentally benign procedure for the oxidation of primary alcohols to the carboxylic acids has been developed. In this procedure, NaClO₂ is used as the stoichiometric oxidant in the presence of catalytic amount of TEMPO

10

and bleach (NaClO). Most primary alcohols were oxidized in essentially quantitative yield. Compared with TEMPO/NaClO/CH₂Cl₂ protocol, the amount of chlorination of electron rich aromatic rings in the substrates were dramatically reduced and the yield and purity of the products improved. Additionally, no chlorinated solvent is used.

The instant invention can be understood further by the following examples, which do not constitute a limitation of the invention.
General:

All substrates and reagents were obtained commercially, except 1g (See Examples 2-5 describing the preparation of this primary alcohol) and used without purification. ¹H and ¹³C NMR spectra were recorded at 250 and 62.5 MHz respectively. The products were identified by comparing their NMR spectra with those of commercial materials except for 2g and 2j. The yields were determined by reverse phase HPLC with Zorbax SB-Phenyl or YMC ODS-AM columns and MeCN/0.1% H₃PO₄ as the mobile phase.

EXAMPLE 1

20 Oxidation of Primary Alcohol—TEMPO Oxidation



A mixture of the primary alcohol 1 (40 mmol) in MeCN (200 mL) and sodium phosphate buffer (0.67 M, pH= 6.7) was heated to 35 °C. TEMPO (436 mg, 2.8 mmol) was added then a solution of sodium chlorite (9.14 g 80%, 80.0 mmol in 40 mL water) and a solution of dilute bleach (1.06 mL 5.25% bleach diluted into 20 mL, 2.0mol%) were added simultaneously in 2 hours.*

*Do not mix the sodium chlorite solution and bleach prior to the addition since the mixture appears to be unstable. The addition should be carried out as follows: approximately 20% of the sodium chlorite solution is added followed by 20%

of the dilute bleach. Then the rest of the NaClO_2 solution and dilute bleach are added simultaneously in 2 hours. The reaction is slightly exothermic.

The mixture was stirred at 35 °C until the reaction is complete (<2A% SM, 2-4 h) then cooled to rt. Water (300 mL) was added and the pH was adjusted to 8.0 with 2.0 N NaOH (~48 mL). The reaction was quenched by pouring into cold (0 °C) Na_2SO_3 solution (12.2 g in 200 mL water) maintained < 20 °C. The pH of the aqueous layer should be 8.5-9.0 After stirring for 0.5 hour at rt, MTBE (200 mL) was added with stirring. The organic layer was separated discarded. More MTBE (300 mL) was added and aqueous layer was acidified with 2.0 N HCl (~100 mL) with stirring to pH = 3-4. The organic layer was washed with water (2 x 100 mL), brine (150 mL) to give a solution of the crude carboxylic acid 2 in 90-95% yield.

EXAMPLE 2

Preparation of 2-bromo-5-methoxybenzyl alcohol



Sodium borohydride (8.6 g) is slurried in THF (150mL KF=150 µg/mL) in a round bottom flask equipped with a thermocouple, an addition funnel, a nitrogen inlet a mechanical stirrer and a cooling bath. 2-Bromo-5-methoxybenzoic acid (50 g) is dissolved in THF (100mL KF= 150 µg/mL) is added to the sodium borohydride slurry over 45 min while maintaining the temperature at 20-25°C. The reaction must be controlled with intermittent cooling and by careful monitoring of the addition rate. The mixture is aged for 30 min at 20-25°C. Boron trifluoride etherate (36.9 g) is added over a period of 30 min at 30-35°C. The addition of boron trifluoride etherate produces a delayed exotherm and should be added slowly in order to control the reaction

temperature. The resulting white slurry is aged for 1 h at 30-35°C and then sampled for HPLC assay. A peak at RT = 8.7 min is an impurity related to the starting material. The acid is at RT = 9.1min.


The reaction mixture is cooled to 15°C and carefully
5 quenched into a cold (10 °C) saturated ammonium chloride solution (150 mL) while maintaining the temperature < 25°C.

Ethyl acetate (500 mL) is added and the layers are separated. The organic layer is washed with water (100 mL) and then transferred to a 1L round bottom flask equipped for distillation. The
10 solution was concentrated and charged with fresh ethyl acetate. This is repeated until a solution with a volume of 200 mL has KF<200 µg/mL. The solvent is then switched to DMF to give the final volume of 200 mL with a KF<200 µg/mL.

15

EXAMPLE 3

Preparation of 2-bromo-5-methoxybenzyl chloride



20

The DMF solution of the benzyl alcohol (91.3 g in 400mL
KF=300 µg/mL) is charged to a 2 L flask equipped with a mechanical stirrer, thermocouple, N₂ inlet, and cooling bath. The solution is cooled to 0-5°C and the addition funnel is charged with thionyl chloride (55.0 g). The thionyl chloride is added over a period of 45 min while maintaining
25 the temperature 5-10°C. The mixture is aged for 1 h at 5°C and assayed by HPLC.


The addition funnel is charged with water (400 mL) which is added dropwise to the reaction mixture over a period of 30 min. while maintaining the temperature < 15°C. The temperature is controlled by

cooling and monitoring the rate of addition. The initial addition of water is highly exothermic. Using large excess of thionyl chloride results in a more exothermic quench. If the quench temperature is not controlled, hydrolysis of the benzyl chloride back to the alcohol may result.

- 5 The resulting thick white slurry is aged for 1 h at 0-5°C. The benzyl chloride is isolated by filtration. The cake is washed with (1:1) DMF:H₂O (100mL) and then water (200 mL). The solid is dried in vacuo to give 93 g of the benzyl chloride(94% yield, 96 A%).
HPLC assay: Column: Waters Symmetry C8, 4.6 x 250mm; UV
10 Detection: 220 nm; Column Temp: 25 °C; Flow rate: 1 mL / min.; Eluent: CH₃CN:H₂O:0.1% H₃PO₄ (70:30); RT (benzyl alcohol) = 3.9 min; RT (benzyl chloride) = 7.3 min.; and RT (DMF) = 2.6 min.

EXAMPLE 4

- 15 Preparation of the Acetonide of N-propanoyl (1R,2S)-cis-aminoindanol



- A 5 L 3-neck round bottom flask equipped with a mechanical stirrer, N₂ inlet, thermocouple probe, heating mantle, and
20 addition funnel is charged with (1R,2S)-cis-aminoindanol (100 g), tetrahydrofuran (1.2 L, KF 120 µg/mL), and triethylamine (96 mL, KF 500 µg/mL). The resulting slurry is heated under a N₂ atmosphere to 40-45°C giving a yellow solution. Propionyl chloride (59 mL) is charged to an addition funnel and added to the solution while maintaining the
25 temperature at 45-50°C.

The temperature is controlled by rate of propionyl chloride addition and a cooling bath. HPLC assay shows >99% amide formed. Methanesulfonic acid (3 mL) is added to the reaction slurry. 2-

Methoxypropene (140 mL) is charged to an addition funnel and added over 30 minutes at a temperature of 50°C.

The addition of 2-methoxypropene is mildly exothermic. The temperature is maintained by the rate of addition and a heating mantle. The reaction remains a slurry but does become less thick.


The reaction slurry is aged for 1-2 hours at 50°C. HPLC assay at this point shows <0.5A% of the amide remaining. The amide is not removed in the isolation so it is important to push the reaction to completion. The reaction slurry is cooled to 0-5°C and quenched by addition of 5% aqueous sodium carbonate solution (1 L) and heptane (1 L). The layers are stirred and separated and the organic is washed with water (300 mL).

HPLC assay at this point shows the acetone in >98A% and >90% yield. The acetone/THF/heptane solution is filtered into a 2 L round bottom flask and the solution is distilled to a final volume of 700mL. Heptane (1L) is added and the solution is distilled to a final volume of 700mL. The distillation is done under partial vacuum at -50°C. NMR assay at this point shows < 2 mol% THF. The solution is allowed to cool and is seeded with acetone at 35-40°C. The thick slurry is aged for 1 hour at ambient temperature then cooled to 0-5°C and aged for 1 hour. The slurry is filtered and the cake is washed with cold heptane (200 mL) and air dried to yield acetone as a crystalline white solid (141.1 g, 85% yield, 99.6 A%).

25

EXAMPLE 5

Alkylation of the Acetone with 2-bromo-5-methoxybenzyl chloride.



A THF solution (2L, KF < 200 µg/mL) of the acetamide (252 g) and the benzyl chloride (255 g) is cooled to -10°C. Lithium bis(trimethylsilyl)amide (1.45 L) is added dropwise over 5 h at 0-2°C. The mixture is then aged for 1.5 h and assayed by HPLC.

5 The reaction is quenched by adding aqueous saturated ammonium chloride solution (1 L). The initial addition of the ammonium chloride should be slow in order to control the foaming. The rate can be increased when the foaming subsides.

10 The quenched reaction is then transferred into a mixture of aqueous ammonium chloride (1.5 L), water (0.5 L), and ethyl acetate (3 L). The mixture is then agitated for 15 min and the layers are separated. The organic layer is washed with water (1 L) and brine (0.5 L). The ethyl acetate solution is concentrated to a low volume and solvent switched to 1,4-dioxane. The dioxane solution is adjusted to a final volume of 1.8 L.

15 The dioxane solution of the coupled product is charged to a 12 L round bottom flask and 6 M HCl (1.5 L) is charged. The mixture is heated to reflux and monitored by HPLC.


20 The mixture is cooled to 20°C and MTBE (3 L) is added. The mixture is agitated for 15 min and the layers are separated. The organic layer is washed with water (1 L). The MTBE solution of the crude acid is extracted with 0.6 M sodium hydroxide (2 L). The aqueous solution of the sodium salt of the acid is combined with MTBE (2.5 L) and cooled to 10°C.

25 The two phase mixture is acidified with 5.4 M sulfuric acid (250 mL), agitated for 15 min, settled and the layers separated. The MTBE solution of the acid is washed with water (0.5 L). The MTBE solution of the acid is dried by distillation and then solvent switched to THF. The final volume of the THF is 2 L with a KF < 250 µg/mL.

30 HPLC assay: column: Waters Symmetry; Eluent: acetonitrile: water: phosphoric acid (70:30:0.1); Flow rate: 1 mL/min.; RT (acetamide) = 4.5 min.; RT (benzyl chloride) = 7.5 min.; RT (coupled product) = 11.5 min.; RT (aminodanol) = 1.7 min.; RT (hydroxyamide) = 1.7 min.; and RT (acid) = 4.5 min.

EXAMPLE 6

Preparation of 3-(2-bromo-5-methoxyphenyl)-2-methylpropanol



Sodium borohydride (33 g) is slurried in THF (0.5 L KF=200
5 $\mu\text{g/mL}$) in a round bottom flask. The THF solution (2 L) of the acid is
added to the sodium borohydride slurry over 1 h while maintaining the
temperature at 20-25°C.

The reaction is controlled with a cooling bath and by
carefully monitoring the addition rate. A nitrogen sweep and proper
10 venting of the hydrogen is also important.

The mixture is aged for 30 min at 20-25 °C. Boron trifluoride
etherate (152 g) is added over 1 h at 30-35 °C. The addition produces a
delayed exotherm and should be carefully monitored in order to control
the reaction temperature. The resulting milky white slurry is aged for 1
15 h at 30 °C and sampled for HPLC assay.


The reaction mixture is cooled to 15 °C and carefully
quenched in a cold (10°C) ammonium chloride solution (1.5 L) while
maintaining the temperature at 25 °C. The rate of hydrogen evolution is
controlled by the rate of the addition of the mixture into the ammonium
20 chloride. The quenched mixture is distilled *in vacuo* to remove the
THF. The aqueous layer is extracted with MTBE (1.5 L) and the organic
layer is dried by flushing with additional MTBE. The MTBE solution is
then solvent switched to hexanes and adjusted to a volume of 350 mL and
seeded. The slurry is aged for 2 h at 20 °C and then cooled to 0-5 °C aged
25 for 1 h and filtered. The cake is washed with cold hexanes (200 mL). The
solid is dried under a nitrogen sweep. The isolated solid (164 g) is >
99A% by HPLC and > 99%ee.

HPLC: Column: Waters Symmetry C8; Solvent: acetonitrile:water: phosphoric acid (50:50:0.1); Flow rate: 1mL /min.; Detection: 220 nm; RT (acid) = 10.2 min.; RT (alcohol) = 10.7min.

5 Chiral HPLC: Column: Chiracel OD-H; Hexane:2-propanol (97:3); Flow rate: 1 mL/ min.; Detection: 220 nm; RT minor isomer = 21 min.; and RT major isomer = 23 min.

EXAMPLE 7

Preparation of 3-(2-bromo-5-methoxyphenyl)-2-methylpropanoic acid



The acid was prepared following the general procedure recited in Example 1.

2g: ^1H NMR (CDCl_3) δ : 7.44 (d, $J=8.7$ Hz, 1H), 6.78 (d, $J=3.1$ Hz, 1H), 6.66 (dd, $J=8.7, 3.1$ Hz, 1H), 3.75 (s, 3H), 3.13 (dd, $J=13.1, 6.8$ Hz, 1H), 2.98-2.84 (m, 1H), 2.77 (dd, $J=13.1, 7.4$ Hz, 1H), 1.23 (d, $J=6.9$ Hz, 3H).


2j: ^1H NMR (CDCl_3) δ : 9.0-8.0 (broad, 1 H), 7.47-7.30 (m, 5H), 5.71 (d, $J = 7.7$ Hz, 1H), 4.43 (d, $J = 7.7$ Hz, 1H), 2.70-2.40 (m, 2H), 2.33-2.27 (m, 1H), 2.17-1.80 (m, 3H), 1.58 (s, 3H).

^{13}C NMR (CDCl_3) δ : 172.04, 169.48, 137.52, 128.73, 126.16, 94.66, 77.05, 64.34, 34.52, 29.91, 23.45, 17.28.

Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NO}_4$: C, 65.44; H, 6.22; N, 5.09. Found C, 65.31; H, 6.15; N, 4.98.

WHAT IS CLAIMED IS:

1. A process for preparing a compound of Formula I:



wherein:

5 R^1 is:

- a) H,
- b) C₁-C₈ alkyl,
- c) C₂-C₈ alkynyl,
- d) C₃-C₇ cycloalkyl,
- 10 e) aryl,
- f) heteroaryl, or
- g) heterocyclyl;

C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl, are
 15 unsubstituted or substituted with one, two or three
 substituents selected from the group consisting of: OH,
 CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₃-C₈ cycloalkyl,
 aryl, heteroaryl, heterocyclyl, and CO(CH₂)_nCH₃,

aryl is defined as phenyl or naphthyl, which is unsubstituted or
 20 substituted with one, two or three substituents selected from
 the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈
 alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, or C₃-C₈ cycloalkyl,
 CO(CH₂)_nCH₃, or when aryl is substituted on adjacent
 carbons they can form a 5- or 6-membered fused ring
 having one, two or three heteroatoms selected from O, N,
 25 and S, this ring is unsubstituted or substituted on carbon or
 nitrogen with one, two or three substituents selected from
 the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈
 alkoxy, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₃-C₈
 cycloalkyl, and CO(CH₂)_nCH₃;

30 heteroaryl is defined as a 5- or 6-membered aromatic ring
 containing 1, 2 or 3 heteroatoms selected from O, N and S,

which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and additionally
5 the 5- or 6-membered aromatic ring can be benzofused and unsubstituted or substituted with one, two or three substituents as described above;

heterocyclyl is defined as a 5- or 6-membered, non-aromatic ring containing 1, 2 or 3 heteroatoms selected from O, N and S ,
10 which may contain one or two double bonds and which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of: OH, CO₂R⁴, Br, Cl, F, I, CF₃, C₁-C₈ alkoxy, C₁-C₈ alkyl, C₂-C₈ alkynyl, C₃-C₈ cycloalkyl, CO(CH₂)_nCH₃, and additionally
15 the 5- or 6-membered ring can be benzofused and unsubstituted or substituted with one, two or three substituents as described above;


20 n is: 0 to 5;

t is: 0, 1 or 2;

R⁴ is: H, or C₁-C₈ alkyl; or

25 comprising the following steps:

- 1) adding to a compound of Formula II in a solvent,



a solution of phosphate buffer to maintain a pH of about 4.0 to about 8.0;

30

- 2) maintaining the phosphate-buffered biphasic mixture of the compound of Formula II at about 0°C to about 50°C;

- 3) adding a catalytic amount of TEMPO to the mixture; and
- 4) charging the TEMPO/phosphate-buffered biphasic mixture —
with a solution of sodium chlorite and a catalytic amount of
sodium hypochlorite to oxidize to the compound of Formula I.

2. The process as recited in Claim 1, wherein the solvent is selected from the group consisting of: acetonitrile, tetrahydrofuran, diethyl ether, MTBE (methyl t-butyl ether), DME (dimethoxyethane), DIGLYME (2-methoxyethyl ether), TRIGLYME (triethylene glycol dimethyl ether), toluene, benzene, hexane, pentane, dioxane, or a mixture of said solvents, including a mixture of said solvents with water.

3. The process as recited in Claim 2, wherein the phosphate buffer comprises an aqueous mixture of NaOH, KOH, NaH_2PO_4 , KH_2PO_4 , Na_2HPO_4 , and K_2HPO_4 , sufficient to maintain a pH of about 4.0 to about 8.0.

4. The process as recited in Claim 3, wherein TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy, free radical) is used in about 1.0 to about 10.0 mole percent.

5. The process as recited in Claim 4, wherein sodium chlorite is used in about 1.0 to about 3.0 equivalents.

6. The process as recited in Claim 5, wherein sodium hypochlorite is used in about 1.0 to about 7.0 mole percent.

7. The process as recited in Claim 6, wherein the reaction temperature is about 0°C to about 50°C.

8. The process as recited in Claim 7, wherein the phosphate buffer comprises an aqueous mixture of NaOH, KOH,

NaH_2PO_4 , KH_2PO_4 , Na_2HPO_4 , and K_2HPO_4 , sufficient to maintain a pH of about 6.5 to about 7.0.

9. The process as recited Claim 8, wherein TEMPO
5 (2,2,6,6-tetramethyl-1-piperidinyloxy, free radical) is used in about 5.0 to about 7.0 mole percent.

10. The process as recited in Claim 9, wherein sodium
chlorite is used in about 2.0 equivalents relative to the compound of
10 Formula II.

11. The process as recited in Claim 10, wherein sodium
hypochlorite is used in about 2.0 to about 5.0 mole percent.

12. The process as recited in Claim 11, wherein the
15 reaction temperature is about 35°C to about 40°C.

13. The process as recited in Claim 12, wherein the
reaction time is about 2 hours to about 4 hours.

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US99/07466

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07C 51/16

US CL : 562/409

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 562/409

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CA OLD, CA PLUS

search terms: CA Reg numbers for TEMPO, NaClO₂, NaClO

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	NOOY, A.E.J. et al. On the use of stable organic radicals for the oxidation of primary and secondary alcohols. synthesis. October 1996, pages 1153-1174, especially pages 1161-1164.	1-13
Y	US 5,631,366 A (LOHRI et al) 20 May 1997, col 3 line 13 - col 4 line 55.	1-13
Y	ANELLI, P.L. et al. Fast and selective oxidation of primary alcohols of aldehydes or to carboxylic acids and of secondary alcohols to ketones mediated by oxoammonium salts under two-phase conditions. J. Org. Chem. 1987, Vol. 52, No. 12, pages 2559-2562, especially page 2562.	1-13

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 JUNE 1999

Date of mailing of the international search report

26 JUL 1999

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT DEEMIE

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/07466

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DALCANALE, E. Selective oxidation of aldehydes to carboxylic acids with sodium chlorite-hydrogen peroxide. J. Org. Chem. 1986, Vol. 51, No. 4, pages 567-571, see entire document.	1-13